



# WOMAN'S RATE OF INFECTION FROM ONE ACT

## OF INTERCOURSE WITH AN INFECTED PARTNER

### # NEW CASES EACH YEAR

#### CURABLE, NONVIRAL DISEASES

female

male

##### **CHLAMYDIA** (4-12 MILLION/YR.)

Can cause PID, Ectopic Pregnancy, Infertility,

↑ risk of HIV if exposed\*, Premature Delivery

---

##### **TRICHOMONIASIS** (3 MILLION/YR)

↑ risk of HIV if exposed\*, Premature Delivery

No data

No data

---

##### **GONORRHEA** (1.1 MILLION/YR)

Can cause PID, Ectopic Pregnancy, Infertility,

↑ risk of HIV if exposed \*, Blinds baby

---

##### **SYPHILIS** (120,000/YR)

Seriously damages many parts of the body,

↑ risk of HIV if exposed \*, Kills/damages baby

---

#### NONCURABLE, VIRAL DISEASES

##### **HUMAN PAPILLOMA VIRUS**

##### **(HPV) GENITAL WARTS** (½ - 1 MILLION/YR)

Some types cause cancer of cervix, vulva, vagina or anus

---

##### **GENITAL HERPES** (200,000-500,000/YR)

Repeated outbreaks, ↑ risk of HIV if exposed \*,

By age 30, 20% Americans infected

Premature Delivery, Kill/damage baby

---

##### **HEPATITIS B** (100,000-200,000/YR)

Promotes liver damage and ↑ risk of liver cancer

---

##### **HIV/AIDS** (40,000-50,000/YR)

HIV infection can progress to AIDS,

Increases risk of other STDs

\*\*Not from one act, Risk of Giving HIV to Partner

**HIV+ Male to HIV - Female**

Woman has an .....risk

of getting infected from the man

---

\* **NOTE** - If exposed to the HIV/AIDS virus, anyone who is infected with these STDs has an increased risk of becoming infected. Anyone infected with the HIV/AIDS virus also has an increased risk of getting infected, if they are exposed to the other STDs.